New Insights into the Sun (NIIS2009): manuscript hg˙niis2009
November 26, 2009

arXiv:0911.5044v1 [astro-ph.SR] 26 Nov 2009

Seismic signature of helium ionization in the Sun and stars

G. Houdek[1] and D.O. Gough[2,3]

[1] Institute of Astronomy, University of Vienna, A-1180 Vienna, Austria, e-mail: `guenter.houdek@univie.ac.at`
[2] Institute of Astronomy, University of Cambridge, Cambridge CB3 0HA, UK, e-mail: `douglas@ast.cam.ac.uk`
[3] Department of Applied Mathematics and Theoretical Physics, University of Cambridge CB3 0WA, UK

ABSTRACT

We present a summary of an asteroseismic signature of helium ionization reported by Houdek & Gough (2007, 2008, 2009) for low-degree p modes in solar-type stars, and illustrate its applications for asteroseismic diagnoses.

Key words. stellar interior - helium abundance - asteroseismology

1. Introduction

Abrupt variation in the stratification of a star (relative to the scale of the inverse radial wavenumber of a seismic mode of oscillation), such as that resulting from the (smooth, albeit acoustically relatively abrupt) depression in the first adiabatic exponent $\gamma_1 = (\partial \ln p/\partial \ln \rho)_s$ caused by the ionization of helium, where p, ρ and s are pressure, density and specific entropy, or from the sharp transition from radiative to convective heat transport at the base of the convection zone, induces small-amplitude oscillatory components (with respect to frequency) in the spacing of the cyclic eigenfrequencies ν_i of seismic modes, with $i = (n, l)$, where n and l are order and degree respectively. We shall call such abrupt variations in the stellar stratification an acoustic glitch. Our interest is principally in the near-surface glitch caused by helium ionization.

2. The seismic diagnostic and discussion

It was demonstrated by Houdek & Gough (2007) that the variation of the sound speed induced by helium ionization enables one to determine from the low-degree eigenfrequencies a measure that is almost proportional to the helium abundance, with little contamination from other properties of the structure of the star. The deviation

$$\delta\nu_i := \nu_i - \nu_{\mathrm{s}i} \tag{1}$$

of the eigenfrequency from the corresponding frequency $\nu_{\mathrm{s}i}$ of a similar smoothly stratified, i.e. glitch-free, star is the indicator of a measure of the helium abundance Y. The individual frequency contributions arising from the acoustic glitches to the total frequency contribution $\delta\nu_i$ are:

$$\delta\nu_i = \delta_\gamma\nu_i + \delta_{\mathrm{c}}\nu_i + \delta_{\mathrm{u}}\nu_i\,, \tag{2}$$

where $\delta_\gamma\nu_i$ and $\delta_{\mathrm{c}}\nu_i$ arise respectively from the variation in γ_1 induced by helium ionization and from the acoustic glitch at the base of the convection zone caused by a near discontinuity in the second derivative of density. Approximate expressions for $\delta_\gamma\nu_i$ and $\delta_{\mathrm{c}}\nu_i$ were presented by Houdek & Gough (2007, 2008). The additional upper-glitch component $\delta_{\mathrm{u}}\nu_i$, which is produced by the ionization of hydrogen and the upper superadiabatic boundary layer of the envelope convection zone, is difficult to model. Houdek & Gough (2007, 2008) estimated all three glitch components by fitting to second differences $\Delta_{2i}\nu := \nu_{n-1,l} - 2\nu_{n,l} + \nu_{n+1,l}$ of the observed frequencies ν_i the seismic diagnostic



Fig. 1. The symbols denote contributions $\delta\nu_i$ to the frequencies ν_i produced by the acoustic glitches of the Sun.

$$\Delta_{2i}\nu \simeq \Delta_{2i}(\delta_\gamma\nu + \delta_{\mathrm{c}}\nu + \delta_{\mathrm{u}}\nu)\,, \tag{3}$$

and represented the upper-glitch component by a series of inverse powers of ν_i, i.e. $\Delta_{2i}\delta_{\mathrm{u}}\nu = \sum_{k=0}^{3} a_k\nu_i^{-k}$, from which the four coefficients a_k, additional to seven more coefficients in $\delta_\gamma\nu_i$ and $\delta_{\mathrm{c}}\nu_i$, were determined. The upper glitch frequency component is represented by (Houdek & Gough 2009)

$$\begin{aligned}\delta_{\mathrm{u}}\nu_i &\simeq A + B\nu_i \\ &+ \left[\frac{1}{2}a_0\nu_i^2 + a_1\nu_i(\ln\nu_i - 1) - a_2\ln\nu_i + \frac{1}{2}a_3\nu_i\right]h^{-2} \\ &\equiv A + B\nu_i + F_{\mathrm{u}}\,,\end{aligned} \tag{4}$$



Fig. 2. The symbols in the **upper panels** denote second differences $\Delta_{2i}\nu := \nu_{n-1,l} - 2\nu_{n,l} + \nu_{n+1,l}$ for low-degree modes obtained from the BiSON (**left panel**; Basu et al. 2007) and from simulated data for a $1\,\mathrm{M}_\odot$ model of age 5.54 Gy (**right panel**); the simulation was based on two periods of 4-months of observation, separated by 1 year, with the SONG network (see Grundahl et al. 2007). The solid curves are fits to $\Delta_{2i}\nu$ based on the analysis by Houdek & Gough (2007). The dashed curves are the smooth contributions, including a third-order polynomial in ν_i^{-1} to represent the upper-glitch contribution from near-surface effects. The **lower panels** display the remaining individual contributions from the acoustic glitches to $\Delta_{2i}\nu$: the dotted and solid curves are the contributions from the first and second stages of helium ionization, and the dot-dashed curve is the contribution from the acoustic glitch at the base of the convective envelope.

where $h = (\nu_{n+1,l} - \nu_{n-1,l})/2$. The somewhat arbitrarily chosen power series in ν_i^{-1} for the upper-glitch component leads to the two undetermined constants of summation A and B, which Houdek & Gough (2009) chose by minimizing

$$E \equiv ||\delta_{\mathrm{u}}\nu_i||_2 = \sum_n (A + B\nu_i + F_{\mathrm{u}})^2 \,. \tag{5}$$

In Fig. 1 is displayed the sum of all acoustic glitches, i.e. Eq.(2), for low-degree solar frequencies observed by BiSON (Basu et al. 2007). Displayed in the top panels of Fig. 2 are the second differences of $\delta\nu_i$ (solid curve) and of the upper-glitch component $\delta_{\mathrm{u}}\nu_i$ (dashed curve) for the Sun and for simulated data of a solar-like star observed with the Danish SONG instrument. The corresponding individual frequency contributions $\delta_\gamma\nu_i$ (dotted and solid curves for the two stages of helium ionization) and $\delta_{\mathrm{c}}\nu_i$ (dot-dashed curve) are illustrated in the lower panels of Fig. 2.

The amplitudes of the oscillatory contributions from the helium ionization (solid and dotted curves in the lower panels of Fig. 2) provide an independent measure of the helium abundance of the star. Moreover, with the help of the smoothed (glitch-free) eigenfrequencies $\nu_{\mathrm{s}i}$ the core-sensitive coefficients of the asymptotic representation of solar-like oscillations (Tassoul 1980, Gough 1986) can be evaluated more accurately. This information stabilizes the age calibration of the Sun and solar-like stars using only low-degree acoustic modes (Houdek & Gough 2008, 2009).

Acknowledgements. Support by the Austrian Science Fund (FWF project P21205) is gratefully acknowledged. Participation at the Ponte de Lima workshop was supported by the European Helio- and Asteroseismology Network (HELAS), a major international collaboration funded by the European Commission's Sixth Framework Programme.

References

Basu S., Chaplin, W. J., Elsworth, Y., New, A. M., Serenelli, G., Verner, G. A. 2007, ApJ, 655, 660

Gough D.O. 1986, in: Proc. Hydrodynamic and magnetohydrodynamic problems in the Sun and stars, ed. Y. Osaki, University of Tokyo, Tokyo, p. 117

Grundahl F., Kjeldsen H., Christensen-Dalsgaard J., Arentoft T., Frandsen S. 2007, CoAst 150, 300

Houdek G., Gough D.O. 2007, MNRAS 375, 861.

Houdek G., Gough D.O. 2008, In The Art of Modelling Stars in the 21st Century, eds Deng, L., Chan, K. L., Chiosi C., IAU Symp., Vol. 252, CUP, Cambridge , p. 149.

Houdek G., Gough D.O. 2009, CoAst 159, 27

Tassoul M. 1980, ApJS 43, 469